

TWMS J. App. Eng. Math. V.7, N.1, 2017, pp. 58-65

ON APPROXIMATE METHODS FOR FRACTAL VEHICULAR TRAFFIC FLOW

H. K. JASSIM¹, §

ABSTRACT. In this paper, we find the approximate solutions for partial differential equations arising in fractal vehicular traffic flow by using the local fractional Laplace decomposition method (LFLDM) and local fractional series expansion method (LFSEM). These methods provide us with a convenient way to find the approximate solution with less computation as compared with local fractional variational iteration method. The results obtained by the proposed methods (LFLDM) and (LFSEM) are compared with the results obtained by (LFLVIM). Some examples are presented to illustrate the efficiency and accuracy of the proposed methods.

Keywords: Lighthill-Whitham-Richards model, local fractional Laplace decomposition method, local fractional series expansion method, fractal vehicular traffic flow.

AMS Subject Classification: 26A33, 35R11, 74H10.

1. INTRODUCTION

Lighthill and Whitham (1955), and Richards (1956) independently presented a macroscopic model of traffic flow to describe the dynamic characteristics of traffic on a homogeneous and unidirectional highway, which is now known as the LWR model in the literature of traffic flow theory. The LWR model is still widely used for the modeling of traffic flow, because of its simplicity and good explanatory power to understand the qualitative behavior of road traffic [1].

The Lighthill-Whitham-Richards model was studied by Li [2] and Wang [3] on a finite length highway and reads as follows

$$\frac{\partial^\vartheta M(x, t)}{\partial t^\vartheta} + \mu \frac{\partial^\vartheta M(x, t)}{\partial x^\vartheta} = 0, \quad 0 < \vartheta \leq 1 \quad (1)$$

where the initial and boundary conditions are presented as follows

$$\begin{aligned} M(x, 0) &= \varphi(x); \\ M(0, t) &= \psi(t). \end{aligned} \quad (2)$$

There are some developed technologies to solve partial differential equations with local fractional operator such as the local fractional variational iteration method [4, 5], local fractional series expansion method [6, 7], local fractional decomposition method [7, 8], local fractional function decomposition method [9, 10], local fractional Laplace variational iteration method [2, 11, 12], and local fractional reduce differential transform method [13].

¹ Department of Mathematics, Faculty of Education for Pure Sciences, University of Thi-Qar, Iraq.
e-mail: hassan.kamil@yahoo.com;

§ Manuscript received: January 1, 2016; accepted: July 15, 2016.

TWMS Journal of Applied and Engineering Mathematics, V.7, No.1; © Işık University, Department of Mathematics, 2017; all rights reserved.

In this work, two methods; namely the local fractional Laplace decomposition method and local fractional series expansion method with local fractional operators, are proposed to solve the linear partial differential equations arising in fractal vehicular traffic flow.

2. LOCAL FRACTIONAL LAPLACE DECOMPOSITION METHOD

Let us consider the following, the local fractional differential operator;

$$L_{\vartheta}M(x, t) + R_{\vartheta}M(x, t) = h(x, t) \quad (3)$$

where $L_{\vartheta} = \frac{\partial^{\vartheta}}{\partial t^{\vartheta}}$ denotes the linear local fractional differential operator, R_{ϑ} is the remaining linear operator, and $h(x, t)$ is a source term.

Taking local fractional Laplace transform on (3), we obtain

$$\tilde{L}_{\vartheta} \{L_{\vartheta}M(x, t)\} + \tilde{L}_{\vartheta} \{R_{\vartheta}M(x, t)\} = \tilde{L}_{\vartheta} \{h(x, t)\}. \quad (4)$$

By applying the local fractional Laplace transform differentiation property, we have

$$s^{\vartheta} \tilde{L}_{\vartheta} \{M(x, t)\} - M(x, 0) + \tilde{L}_{\vartheta} \{R_{\vartheta}M(x, t)\} = \tilde{L}_{\vartheta} \{h(x, t)\}. \quad (5)$$

or equivalently

$$\tilde{L}_{\vartheta} \{M(x, t)\} = \frac{1}{s^{\vartheta}} M(x, 0) + \frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{h(x, t)\} - \frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{R_{\vartheta}M(x, t)\}. \quad (6)$$

Taking the inverse of local fractional Laplace transform on Eq. (6), we have

$$M(x, t) = M(x, 0) + \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{h(x, t)\} \right] - \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{R_{\vartheta}M(x, t)\} \right]. \quad (7)$$

We are going to represent the solution in an infinite series given below:

$$M(x, t) = \sum_{n=0}^{\infty} M_n(x, t). \quad (8)$$

Substituting (8) into (7), which give us this result:

$$\sum_{n=0}^{\infty} M(x, t) = M(x, 0) + \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{h(x, t)\} \right] - \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ R_{\vartheta} \sum_{n=0}^{\infty} M_n(x, t) \right\} \right]. \quad (9)$$

When we compare the left and right hand sides of (9) we obtain

$$\begin{aligned} M_0(x, t) &= M(x, 0) + \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{h(x, t)\} \right], \\ M_1(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{R_{\vartheta}M_0(x, t)\} \right], \\ M_2(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{R_{\vartheta}M_1(x, t)\} \right], \\ M_3(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \{R_{\vartheta}M_2(x, t)\} \right], \\ &\vdots \end{aligned} \quad (10)$$

The recursive relation, in general form is

$$\begin{aligned} M_0(x, t) &= M(x, 0) + \tilde{L}^{-1} \left[\frac{1}{s^\vartheta} \tilde{L}_\vartheta \{h(x, t)\} \right], \\ M_{n+1}(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{1}{s^\vartheta} \tilde{L}_\vartheta \{R_\vartheta M_n(x, t)\} \right]. \end{aligned} \quad (11)$$

3. LOCAL FRACTIONAL SERIES EXPANSION METHOD

We can written the equation (1) in the form

$$\frac{\partial^\vartheta M(x, t)}{\partial t^\vartheta} = R_\vartheta M(x, t), \quad (12)$$

where $R_\vartheta M(x, t) = -\mu L_x^{(\vartheta)} M(x, t)$ is linear local fractional derivative operator of order ϑ with respect to x .

In accordance with the results in [6, 7], there are multiterm separated functions of independent variables x and t reads as

$$M(x, t) = \sum_{i=0}^{\infty} T_i(t) \Phi_i(x), \quad (13)$$

where $T_i(t)$ and $\Phi_i(x)$ are local fractional continuous functions.

In view of (13), we consider

$$T_i(t) = \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)}, \quad (14)$$

so that

$$M(x, t) = \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} \Phi_i(x). \quad (15)$$

In view of (15), we obtain

$$\frac{\partial^\vartheta M(x, t)}{\partial t^\vartheta} = \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} \Phi_{i+1}(x), \quad (16)$$

and

$$R_\vartheta M(x, t) = R_\vartheta \left(\sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} \Phi_i(x) \right) = \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} (R_\vartheta \Phi_i)(x), \quad (17)$$

Making use of (17), we get

$$\sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} \Phi_{i+1}(x) = \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} (R_\vartheta \Phi_i)(x). \quad (18)$$

Hence, from (18), the recursion reads as follows:

$$\Phi_{i+1}(x) = (R_\vartheta \Phi_i)(x). \quad (19)$$

By using (19), we arrive at the following result:

$$M(x, t) = \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1 + i\vartheta)} \Phi_i(x). \quad (20)$$

4. ILLUSTRATIVE EXAMPLES

In this section, we present the initial and boundary value problems for linear partial differential equations arising in fractal vehicular traffic flow.

Example 4.1. *Let us consider the Lighthill-Whitham-Richards model on a finite length highway as*

$$\frac{\partial^\vartheta M(x, t)}{\partial t^\vartheta} + \mu \frac{\partial^\vartheta M(x, t)}{\partial x^\vartheta} = 0, \quad 0 < \vartheta \leq 1 \quad (21)$$

where the initial and boundary conditions are presented as follows:

$$\begin{aligned} M(x, 0) &= E_\vartheta(x^\vartheta); \\ M(0, t) &= \cosh_\vartheta(\mu t^\vartheta) - \sinh_\vartheta(\mu t^\vartheta). \end{aligned} \quad (22)$$

I. By using LFLDM

In view of (11) and (21) the local fractional iteration algorithm can be written as follows:

$$\begin{aligned} M_0(x, t) &= E_\vartheta(x^\vartheta); \\ M_{n+1}(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^\vartheta} \tilde{L}_\vartheta \left\{ \frac{\partial^\vartheta M_n(x, t)}{\partial x^\vartheta} \right\} \right], \quad n \geq 0. \end{aligned} \quad (23)$$

Therefore, from (23) we give the components as follows:

$$M_0(x, t) = E_\vartheta(x^\vartheta); \quad (24)$$

$$\begin{aligned} M_1(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^\vartheta} \tilde{L}_\vartheta \left\{ \frac{\partial^\vartheta M_0(x, t)}{\partial x^\vartheta} \right\} \right] \\ &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^{2\vartheta}} E_\vartheta(x^\vartheta) \right] \\ &= -\frac{\mu t^\vartheta}{\Gamma(1 + \vartheta)} E_\vartheta(x^\vartheta); \end{aligned} \quad (25)$$

$$\begin{aligned} M_2(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^\vartheta} \tilde{L}_\vartheta \left\{ \frac{\partial^\vartheta M_1(x, t)}{\partial x^\vartheta} \right\} \right] \\ &= \tilde{L}_\vartheta^{-1} \left[\frac{\mu^2}{s^{3\vartheta}} E_\vartheta(x^\vartheta) \right] \\ &= \frac{\mu^2 t^{2\vartheta}}{\Gamma(1 + 2\vartheta)} E_\vartheta(x^\vartheta); \end{aligned} \quad (26)$$

$$\begin{aligned} M_3(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^\vartheta} \tilde{L}_\vartheta \left\{ \frac{\partial^\vartheta M_2(x, t)}{\partial x^\vartheta} \right\} \right] \\ &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu^3}{s^{4\vartheta}} E_\vartheta(x^\vartheta) \right] \\ &= -\frac{\mu^3 t^{3\vartheta}}{\Gamma(1 + 3\vartheta)} E_\vartheta(x^\vartheta); \end{aligned} \quad (27)$$

$$\begin{aligned} M_4(x, t) &= -\tilde{L}_\vartheta^{-1} \left[\frac{\mu}{s^\vartheta} \tilde{L}_\vartheta \left\{ \frac{\partial^\vartheta M_3(x, t)}{\partial x^\vartheta} \right\} \right] \\ &= \tilde{L}_\vartheta^{-1} \left[\frac{\mu^4}{s^{5\vartheta}} E_\vartheta(x^\vartheta) \right] \\ &= \frac{\mu^4 t^{4\vartheta}}{\Gamma(1 + 4\vartheta)} E_\vartheta(x^\vartheta). \end{aligned} \quad (28)$$

Consequently, we obtain

$$\begin{aligned} M(x, t) &= E_{\vartheta}(x^{\vartheta}) \left[1 + \frac{\mu^2 t^{2\vartheta}}{\Gamma(1+2\vartheta)} + \frac{\mu^4 t^{4\vartheta}}{\Gamma(1+4\vartheta)} + \cdots \right] \\ &\quad - E_{\vartheta}(x^{\vartheta}) \left[\frac{\mu t^{\vartheta}}{\Gamma(1+\vartheta)} + \frac{\mu^3 t^{3\vartheta}}{\Gamma(1+3\vartheta)} + \cdots \right] \\ &= E_{\vartheta}(x^{\vartheta}) \cosh_{\vartheta}(\mu t^{\vartheta}) - E_{\vartheta}(x^{\vartheta}) \sinh_{\vartheta}(\mu t^{\vartheta}) \end{aligned} \quad (29)$$

II. By using LFSEM

From (19), we obtain the following iterative formula;

$$\Phi_{i+1}(x) = -\mu L_x^{(\vartheta)} \Phi_i(x), \quad (30)$$

where

$$\Phi_0(x) = E_{\vartheta}(x^{\vartheta}). \quad (31)$$

In view of (30) and (31), we obtain the approximations given by

$$\begin{aligned} \Phi_1(x) &= -\mu L_x^{(\vartheta)} \Phi_0(x) = -\mu E_{\vartheta}(x^{\vartheta}); \\ \Phi_2(x) &= -\mu L_x^{(\vartheta)} \Phi_1(x) = \mu^2 E_{\vartheta}(x^{\vartheta}); \\ \Phi_3(x) &= -\mu L_x^{(\vartheta)} \Phi_2(x) = -\mu^3 E_{\vartheta}(x^{\vartheta}); \\ \Phi_4(x) &= -\mu L_x^{(\vartheta)} \Phi_3(x) = \mu^4 E_{\vartheta}(x^{\vartheta}); \\ \Phi_5(x) &= -\mu L_x^{(\vartheta)} \Phi_4(x) = -\mu^5 E_{\vartheta}(x^{\vartheta}); \\ &\vdots \end{aligned} \quad (32)$$

Therefore, by using (20) and (33) we get the solution

$$\begin{aligned} M(x, t) &= \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1+i\vartheta)} \Phi_i(x) \\ &= E_{\vartheta}(x^{\vartheta}) \left[1 - \frac{\mu t^{\vartheta}}{\Gamma(1+\vartheta)} + \frac{\mu^2 t^{2\vartheta}}{\Gamma(1+2\vartheta)} - \frac{\mu^3 t^{3\vartheta}}{\Gamma(1+3\vartheta)} + \frac{\mu^4 t^{4\vartheta}}{\Gamma(1+4\vartheta)} - \cdots \right] \\ &= E_{\vartheta}(x^{\vartheta}) \left[1 + \frac{\mu^2 t^{2\vartheta}}{\Gamma(1+2\vartheta)} + \frac{\mu^4 t^{4\vartheta}}{\Gamma(1+4\vartheta)} + \cdots \right] \\ &\quad - E_{\vartheta}(x^{\vartheta}) \left[\frac{\mu t^{\vartheta}}{\Gamma(1+\vartheta)} + \frac{\mu^3 t^{3\vartheta}}{\Gamma(1+3\vartheta)} + \cdots \right] \\ &= E_{\vartheta}(x^{\vartheta}) \cosh_{\vartheta}(\mu t^{\vartheta}) - E_{\vartheta}(x^{\vartheta}) \sinh_{\vartheta}(\mu t^{\vartheta}). \end{aligned} \quad (33)$$

From Eqs. (29) and (33), the approximate solution of the given problem (21), by using local fractional Laplace decomposition method and local fractional series expansion method, is the same results as that obtained by the local fractional Laplace variational iteration method.

Example 4.2. Let us consider the Lighthill-Whitham-Richards model on a finite length highway as:

$$\frac{\partial^{\vartheta} M(x, t)}{\partial t^{\vartheta}} + \frac{\partial^{\vartheta} M(x, t)}{\partial x^{\vartheta}} = 0, \quad 0 < \vartheta \leq 1 \quad (34)$$

where the initial and boundary conditions are presented as follows:

$$\begin{aligned} M(x, 0) &= \sinh_{\vartheta}(x^{\vartheta}); \\ M(0, t) &= 0. \end{aligned} \quad (35)$$

I. By using LFLDM

In view of (11) and (34) the local fractional iteration algorithm can be written as follows:

$$\begin{aligned} M_0(x, t) &= \sinh_{\vartheta}(x^{\vartheta}); \\ M_{n+1}(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ \frac{\partial^{\vartheta} M_n(x, t)}{\partial x^{\vartheta}} \right\} \right], n \geq 0. \end{aligned} \quad (36)$$

Therefore, from (36) we give the components as follows:

$$M_0(x, t) = \sinh_{\vartheta}(x^{\vartheta}); \quad (37)$$

$$\begin{aligned} M_1(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ \frac{\partial^{\vartheta} M_0(x, t)}{\partial x^{\vartheta}} \right\} \right] \\ &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{2\vartheta}} \sinh_{\vartheta}(x^{\vartheta}) \right] \\ &= -\frac{t^{\vartheta}}{\Gamma(1 + \vartheta)} \sinh_{\vartheta}(x^{\vartheta}); \end{aligned} \quad (38)$$

$$\begin{aligned} M_2(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ \frac{\partial^{\vartheta} M_1(x, t)}{\partial x^{\vartheta}} \right\} \right] \\ &= \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{3\vartheta}} \sinh_{\vartheta}(x^{\vartheta}) \right] \\ &= \frac{t^{2\vartheta}}{\Gamma(1 + 2\vartheta)} \sinh_{\vartheta}(x^{\vartheta}); \end{aligned} \quad (39)$$

$$\begin{aligned} M_3(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ \frac{\partial^{\vartheta} M_2(x, t)}{\partial x^{\vartheta}} \right\} \right] \\ &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{4\vartheta}} \sinh_{\vartheta}(x^{\vartheta}) \right] \\ &= -\frac{t^{3\vartheta}}{\Gamma(1 + 3\vartheta)} \sinh_{\vartheta}(x^{\vartheta}); \end{aligned} \quad (40)$$

$$\begin{aligned} M_4(x, t) &= -\tilde{L}_{\vartheta}^{-1} \left[\frac{\mu}{s^{\vartheta}} \tilde{L}_{\vartheta} \left\{ \frac{\partial^{\vartheta} M_3(x, t)}{\partial x^{\vartheta}} \right\} \right] \\ &= \tilde{L}_{\vartheta}^{-1} \left[\frac{1}{s^{5\vartheta}} \sinh_{\vartheta}(x^{\vartheta}) \right] \\ &= \frac{t^{4\vartheta}}{\Gamma(1 + 4\vartheta)} \sinh_{\vartheta}(x^{\vartheta}). \end{aligned} \quad (41)$$

Consequently, we obtain

$$\begin{aligned} M(x, t) &= \sinh_{\vartheta}(x^{\vartheta}) \left[1 + \frac{t^{2\vartheta}}{\Gamma(1 + 2\vartheta)} + \frac{t^{4\vartheta}}{\Gamma(1 + 4\vartheta)} + \cdots \right] \\ &\quad - \sinh_{\vartheta}(x^{\vartheta}) \left[\frac{t^{\vartheta}}{\Gamma(1 + \vartheta)} + \frac{t^{3\vartheta}}{\Gamma(1 + 3\vartheta)} + \cdots \right] \\ &= \sinh_{\vartheta}(x^{\vartheta}) \cosh_{\vartheta}(t^{\vartheta}) - \sinh_{\vartheta}(x^{\vartheta}) \sinh_{\vartheta}(t^{\vartheta}). \end{aligned} \quad (42)$$

II. By using LFSEM

From (19), we obtain the following iterative formula:

$$\Phi_{i+1}(x) = -L_x^{(\vartheta)} \Phi_i(x), \quad (43)$$

where

$$\Phi_0(x) = \sinh_{\vartheta}(x^{\vartheta}). \quad (44)$$

In view of (43) and (44), we obtain the approximations given by

$$\begin{aligned} \Phi_1(x) &= -L_x^{(\vartheta)}\Phi_0(x) = -\sinh_{\vartheta}(x^{\vartheta}); \\ \Phi_2(x) &= -L_x^{(\vartheta)}\Phi_1(x) = \sinh_{\vartheta}(x^{\vartheta}); \\ \Phi_3(x) &= -L_x^{(\vartheta)}\Phi_2(x) = -\sinh_{\vartheta}(x^{\vartheta}); \\ \Phi_4(x) &= -L_x^{(\vartheta)}\Phi_3(x) = \sinh_{\vartheta}(x^{\vartheta}); \\ \Phi_5(x) &= -L_x^{(\vartheta)}\Phi_4(x) = -\sinh_{\vartheta}(x^{\vartheta}). \\ &\vdots \end{aligned} \quad (45)$$

Therefore, by using (20) and (46) we get the solution

$$\begin{aligned} M(x, t) &= \sum_{i=0}^{\infty} \frac{t^{i\vartheta}}{\Gamma(1+i\vartheta)} \Phi_i(x) \\ &= \sinh_{\vartheta}(x^{\vartheta}) \left[1 - \frac{t^{\vartheta}}{\Gamma(1+\vartheta)} + \frac{t^{2\vartheta}}{\Gamma(1+2\vartheta)} - \frac{t^{3\vartheta}}{\Gamma(1+3\vartheta)} + \frac{t^{4\vartheta}}{\Gamma(1+4\vartheta)} - \cdots \right] \\ &= \sinh_{\vartheta}(x^{\vartheta}) \left[1 + \frac{t^{2\vartheta}}{\Gamma(1+2\vartheta)} + \frac{t^{4\vartheta}}{\Gamma(1+4\vartheta)} + \cdots \right] \\ &\quad - \sinh_{\vartheta}(x^{\vartheta}) \left[\frac{t^{\vartheta}}{\Gamma(1+\vartheta)} + \frac{t^{3\vartheta}}{\Gamma(1+3\vartheta)} + \cdots \right] \\ &= \sinh_{\vartheta}(x^{\vartheta}) \cosh_{\vartheta}(t^{\vartheta}) - \sinh_{\vartheta}(x^{\vartheta}) \sinh_{\vartheta}(t^{\vartheta}). \end{aligned} \quad (46)$$

From Eqs.(42) and (46), the approximate solution of the given problem (34), by using local fractional Laplace decomposition method and local fractional series expansion method, is the same results as that obtained by the local fractional Laplace variational iteration method.

5. CONCLUSIONS

In this work, we have successfully provided new applications of the local fractional Adomian decomposition method and the local fractional series expansion method for solving linear partial differential equations arising in fractal vehicular traffic flow. The approximate solutions are obtained in series form that rapidly converges in a closed exact formula with simply computable terms. These techniques can be extended to solve various linear and nonlinear fractional problems in applied science.

REFERENCES

- [1] Ansorge,R., (1990), What does the entropy condition mean in traffic flow theory, Transportation Research Part B, 24(2), pp.133-143.
- [2] Li,Y., Wang,L.F., Zeng,S D. and Zhao,Y., (2014), Local Fractional Laplace Variational Iteration Method for Fractal Vehicular Traffic Flow, Advances in Mathematical Physics, 2014, pp.1-7.
- [3] Wang,L.F., Yang,X.J., Baleanu,D., Cattani,C. and Zhao,Y., (2014), Fractal Dynamical Model of Vehicular Traffic Flow within the Local Fractional Conservation Laws, 2014, pp.1-5.
- [4] Yang,X.J. and Baleanu,D., (2013), Fractal heat conduction problem solved by local fractional variational iteration method, Thermal Science, 17(2), pp.625-628.
- [5] Xu,S., Ling,X., Zhao,Y. and Jassim,H.K., (2015), A Novel Schedule for Solving the Two-Dimensional Diffusion in Fractal Heat Transfer, Thermal Science, 19(1), pp.99-103.

- [6] Yang,A.M., Yang,X.J. and Li,Z.B., (2013), Local Fractional series expansion method for Solving wave and diffusion Equations on Cantor sets, Abstract and Applied Analysis, 2013, pp.1-6.
- [7] Jafari,H. and Jassim,H.K., (2015), Application of the Local fractional Adomian Decomposition and Series Expansion Methods for Solving Telegraph Equation on Cantor Sets Involving Local Fractional Derivative Operators, Journal of Zankoy Sulaimani-Part A, 17(2), pp.15-22.
- [8] Jafari,H. and Jassim,H.K., (2014), Local Fractional Adomian Decomposition Method for Solving Two Dimensional Heat conduction Equations within Local Fractional Operators, Journal of Advance in Mathematics, 9(4), pp.2574-2582.
- [9] Wang,S.Q., Yang,Y.J. and Jassim,H.K., (2014), Local Fractional Function Decomposition Method for Solving Inhomogeneous Wave Equations with Local Fractional Derivative, Abstract and Applied Analysis, 2014, pp.1-7.
- [10] Yan,S.P., Jafari,H. and Jassim,H.K., (2014), Local Fractional Adomian Decomposition and Function Decomposition Methods for Solving Laplace Equation within Local Fractional Operators, Advances in Mathematical Physics, 2014, pp.1-7.
- [11] Jafari,H. and Jassim,H.K., (2014), Local Fractional Laplace Variational Iteration Method for Solving Nonlinear Partial Differential Equations on Cantor Sets within Local Fractional Operators, Journal of Zankoy Sulaimani-Part A, 16(4), pp.49-57.
- [12] Jassim, H. K., Unlu, C., Moshokoa, S. P. and Khalique, C. M., (2015), Local Fractional Laplace Variational Iteration Method for Solving Diffusion and Wave Equations on Cantor Sets within Local Fractional Operators, Mathematical Problems in Engineering, 2015, pp. 1-9.
- [13] Jafari,H. and Jassim,H.K., Moshokoa,S.P., Ariyan,V.M. and Tchier,F., (2016), Reduced differential transform method for partial differential equations within local fractional derivative operators, Advances in Mechanical Engineering, 8(4), pp.1-6.
- [14] Jafari,H. and Jassim,H.K., (2015), Numerical Solutions of Telegraph and Laplace Equations on Cantor Sets Using Local Fractional Laplace Decomposition Method , International Journal of Advances in Applied Mathematics and Mechanics, 2(3), pp.144-151.



Hassan Kamil Jassim is an assistant professor in the Department of Mathematics Faculty of Education for Pure Sciences at Thi-Qar University, Iraq. He was born in 1980. His research areas are applied mathematics and mathematical physics including the analytical and numerical methods for ordinary and partial differential equations within local fractional derivative operators.
